

ECTO & ENDO INOCULANT  
**DIEHARD™** **Injectable**



**“The Industry's First Complete Injectable Inoculant”**

**Product Features**

**Endomycorrhizal Fungi**  
**Beneficial Bacteria**

**Humic Acid**  
**Sea Kelp**

**Amino Acids**

**Horta-Sorb® Water Management Gel**

**Trichoderma**

**Yucca Plant Extracts**

**Vitamin B Complex**

**Lasting Residual**

**Convenient Pre-Measured Bags**

**Product Benefits**

**IMPROVES**

**Survival**  
**Rooting**  
**Flowering**  
**Water Absorption**  
**Nutrient Availability**  
**Yields and Production**  
**Client Satisfaction**  
**& Goodwill**

**REDUCES**

**Plant Losses**  
**Fertilizer Use**  
**Need Of Pesticides**  
**Heat Stress Damage**  
**Irrigation Frequency**  
**Losses From Drought**  
**Conditions**

**Compatibility**

**Species:** All plants that associate with endo- and ectomycorrhizal fungi.

**Fungicides:** Non-systemic fungicides normally have no effect. Foliar applied fungicides normally have no effect. Systemic fungicides may be applied 2 weeks before or after use of product. Fungicide use according to label instructions do not extinguish mycorrhiza, they only inhibit development temporarily.

**Non Plant Food Ingredients**

<i>Endomycorrhizal Fungi</i>	278 Propagules per gram <i>Glomus mosseae</i> , <i>Glomus intraradices</i> , <i>Glomus claroideum</i> , <i>Glomus etunicatum</i> , <i>Glomus clarum</i> , <i>Glomus deserticola</i> , <i>Glomus microaggregatum</i> .
<i>Ectomycorrhizal Fungi</i>	14,537,445 spores per gram to include: <i>Pisolithus tinctorius</i> (12,502,203), <i>Rhizopogon</i> (2,035,242).
<i>Trichoderma</i>	1,599,119 CFU'S per gram <i>Genus Trichoderma</i> : (6 strains, 2 each) <i>Trichoderma hamatum</i> , <i>Trichoderma harzianum</i> , <i>Trichoderma reesei</i> .
<i>Bacillus, Pseudomonas &amp; Streptomyces</i> Species	464,317,181 CFU's per gram <b>Genus Bacillus:</b> <i>B. megaterium</i> , <i>B. subtilis</i> , <i>B. amyloliquefaciens</i> , <i>B. azotoformans</i> , <i>B. licheniformis</i> , <i>B. pumilus</i> , <i>B. Firmus</i> , <i>B. coagulans</i> , <i>Geobacillus stearothermophilus</i> , and <i>Paenibacillus polymyxa</i> , <i>Paenibacillus Durum</i> . <b>Genus Pseudomonas:</b> <i>P. putida</i> , <i>P. fluorescense</i> , <i>P. Aureofaceans</i> . <b>Genus Streptomyces:</b> <i>S. lydicus</i> , <i>S. griseus</i> , <i>S. coelicolor</i> .

**OTHER INGREDIENTS**

Root Promoting Vitamins	B, B2, B3, B6, B7, B12, C, K, Biotin, Fulvic Acid
Amino Acids	<i>L-Alanine, L-Argine, L-Aspartic Acid, L-Cystine, L-Glutamic Acid, L-Glycine, L-Histidine, L-Isoleucine, L-Leucine, L-Lysine, L-Methionine, L-Phenylalanine, L-Proline, L-Threonine, L-Serine, L-Tryptophan, L-Tyrosine, L-Valine.</i>
Soluble Sea Kelp	13.0% <i>Ascophyllum nodosum</i>
Humic Acids	20.0% Derived from Leonardite
Sol. Yucca Extract	6.0% <i>Yucca schidigera</i>


**Directions For Use**

Mix with 100 gallons of water. Decant, if necessary.

**Established Trees:**

- 1. Root Injection:** Inject in a 2 1/2 to 3 foot grid pattern from the trunk out to the drip line and beyond if possible. Apply 1-2 quarts of solution per injection site. Use up solution by the end of the day.
- 2. Drip Tape and Microjet:** Decant product before adding to mix tank. Apply at the rate of one pound per acre every six months supplemented every 3 months with BioRush® at two pounds per acre. Irrigate after application.

**New Tree Plantings:** Mix with water and use as a mud-in drench when planting new trees. One pound treats 175 one-year trees. Mix with water used to transplant accordingly. **Small and newly planted trees and shrubs:** Inject along the original root ball in 2-3 places using minimum 1/2 gallon for recently planted trees.



ECTO & ENDO INOCULANT

# DIEHARD™ Injectable



**Nursery Propagation Drench:** Each 8 oz. bag treats 5,000 seedlings. Best application method is a 2-step process. Apply 1-2 weeks after sowing just as roots have developed to the sides of cells. Mix with sufficient water to apply as a drench with a backpack sprayer. Care should be taken to ensure that spores applied are drenched into the media to the level of the roots. Second application just as you are hardening off the crop for shipment. One week before and one week after avoid use of phosphate fertilizers. Avoid the use of high phosphorus fertilizer as this treatment would have a negative effect on Endomycorrhizal colonization. Organic nitrogen fertilizers are recommended.

### Plant Species and Type of Mycorrhizae

<u>Endomycorrhizae</u>			
Arborvitae	Cedar	Ginkgo	Morning Glory
Acacia	Celery	Grapes (all)	Rose
Agapanthus	Cherry	Grass	Mulberry
Ailanthus	Chokeberry	Gum	Monkey Pod
Alder	Chrysanthemum	Hackberry	Nasturtium
Alfalfa	Citrus (all)	Hawthorn	Okra
Almond	Clover	Hibiscus	Olive
Apple	Coconut	Holly	Onion
Apricot	Coffee	Hophornbeam	Pacific Yew
Artichoke	Coral Tree	Hombeam	Palms (all)
Ash	Corn	Horsechestnut	Palmetto
Asparagus	Cotton	Impatiens	Pampas Grass
Avocado	Cottonwood	Jojoba	Papaya
Bamboo	Crabapple	Juniper	Passion Fruit
Basil	Cryptomeria	Kiwi	Paulownia
Bayberry	Cucumber	Leek	Paw Paw
Bean	Currant	Lettuce	Pea
Begonia	Cypress	Lily	Peach
Black Locust	Dogwood	Locust	Peanut
Blackberry	Eggplant	London	Pear
Box Elder	Elm	Planetree	Pecan
Buckeye	Eucalyptus	Macadamia	Pepper
Bulbs	Euonymus	Magnolia	Pistachio
Burning Bush	Fern	Mahogany	Persimmon
Cacao	Fescue	Mahonia	Pittosporum
Cactus	Fig	Mango	Plum
Camellia	Forsythia	Maples (all)	Poinsettia
Carrot	Fountain Grass	Marigold	Potato
Casuarina	Fuchsia	Melons (all)	Poplar
Cassava	Gardenia	Mesquite	Raintree
Catalpa	Garlic	Millet	Raphiolepis
Ceanothus	Geranium	Mimosa	Raspberry
			Redbud

All nut trees except Pecan, Chestnut, and Hazelnut.  
 All fruit trees, grapevines, grasses and many vegetables  
 All berries except blueberry, cranberry and lingonberry.  
 All shrubs and foliage except Laurel, Rhododendron, and Azalea

<u>Ectomycorrhizae</u>				
Alder	Birch	Hemlock	Madrone	Poplar
Aspen	Chestnut	Hickory	Oak	Spruce
Basswood	Chinquapin	Larch	Pecan	Willow
Beech	Fir	Linden	Pine	

<u>Both Endo and Ectomycorrhizae</u>		
Alder	Cottonwood	Poplar
Aspen	Eucalyptus	Willow

## Health & Safety

- Inhalation** If dusty conditions exist, wear a dust mask.
- Ingestion** Drink large amounts of water, induce vomiting, and seek medical attention.
- Eyes** Flush with large amounts of water.
- Safety** Product becomes very slippery when wet.

## Storage & Handling

Store in a cool, dry place. Avoid high temperatures and direct sunlight. Product shelf life is up to 24 months.

**Limited Warranty:** All information contained herein is offered in good faith. There are no warranties of merchantability of fitness for a particular purpose which extend beyond the information contained herein. Our liability is limited to replacement of any product which does not meet these specifications.

**Distributed by:**